

# Factors in Polio Vaccine Hesitancy: A Parental Survey

Javeria Ishaq<sup>1</sup>

<sup>1</sup> Hameed Latif Hospital, Lahore, Pakistan

\* Corresponding author: [javeria748@gmail.com](mailto:javeria748@gmail.com)

## ABSTRACT

**Background:** Despite ongoing nationwide immunization campaigns and widespread availability of vaccination services, polio eradication efforts in Pakistan continue to be hindered by persistent vaccine hesitancy. Parental mistrust, misinformation, and sociocultural beliefs remain major contributors to reduced vaccine uptake, particularly in high-risk groups. **Objective:** This study aimed to assess the prevalence of polio vaccine hesitancy among parents visiting a tertiary pediatric hospital in Lahore and to identify independent factors associated with hesitancy. **Methods:** A cross-sectional study was conducted at the Children's Hospital Lahore involving 257 parents or primary caregivers of children aged five years or younger. Data were collected through a structured interviewer-administered questionnaire covering demographic information, polio-related knowledge, perceived vaccine safety, exposure to misinformation, and attitudinal barriers. Analysis included descriptive statistics, chi-square testing, and multivariable logistic regression. **Results:** Among the 257 participants, 71 (27.6%) exhibited polio vaccine hesitancy. Significant factors associated with hesitancy included low educational attainment, low household income, distrust in vaccine safety, belief in infertility myths, religious apprehensions, and exposure to vaccine-related misinformation on social media. In multivariable analysis, distrust in vaccine safety emerged as the strongest predictor of hesitancy (adjusted OR 5.72; 95% CI 2.76–11.87), followed by belief in infertility myths (adjusted OR 4.38; 95% CI 2.01–9.53) and exposure to misinformation (adjusted OR 3.16; 95% CI 1.68–5.93). **Conclusion:** Polio vaccine hesitancy among parents in Lahore remains substantial and is largely driven by mistrust and misinformation. Strengthening vaccine acceptance will require focused communication strategies, targeted myth-busting approaches, and trust-building interventions. **Keywords:** polio; vaccine hesitancy; parents; Lahore; misinformation; trust; Pakistan

**"Cite this Article"** | Received: 18 March 2025; Accepted: 21 September 2025; Published: 30 September 2025.

**Author Contributions:** Concept: JI; Design: JI; Data Collection: JI; Analysis: JI; Drafting: JI. **Ethical Approval:** Hameed Latif Hospital, Lahore. **Informed Consent:** Written informed consent was obtained from all participants; **Conflict of Interest:** The authors declare no conflict of interest; **Funding:** No external funding; **Data Availability:** Available from the corresponding author on reasonable request; **Acknowledgments:** N/A.

## INTRODUCTION

Poliomyelitis remains a major vaccine-preventable illness targeted for global eradication through coordinated immunization initiatives. While the Global Polio Eradication Initiative has eliminated wild poliovirus from most regions worldwide, Pakistan and Afghanistan continue to experience endemic transmission, creating ongoing challenges for international public health efforts. Despite extensive supplementary immunization campaigns—such as the Expanded Program on Immunization (EPI) and National Immunization Days—persistent pockets of under-immunized communities still jeopardize eradication progress. In this context, vaccine hesitancy, defined as delaying or refusing vaccination despite the availability of services, has emerged as a key barrier to achieving sufficient population immunity.

Growing evidence shows that parental perceptions and attitudes strongly influence childhood vaccination uptake. Community mistrust, misinformation, and sociocultural beliefs are repeatedly identified as major drivers of vaccine refusal in low- and middle-income countries. In Pakistan,

misconceptions about vaccine safety, fears of infertility, and distrust toward vaccination teams have contributed to reduced immunization coverage. Additionally, misinformation circulated through informal networks and social media has intensified skepticism and weakened public confidence in vaccination efforts.

Studies from various regions of Pakistan consistently report high levels of polio vaccine hesitancy among parents. Research in high-risk areas of Karachi has highlighted concerns about vaccine quality and possible side effects as leading reasons for refusal. Findings from Mardan further show that socioeconomic challenges, low literacy, and rural residence are linked to reduced acceptance of polio immunization. Investigations from Khyber Pakhtunkhwa and other areas also emphasize that parental misconceptions and limited knowledge about vaccine-preventable diseases significantly shape vaccination behavior. Together, these findings demonstrate that vaccine hesitancy in Pakistan is deeply rooted in sociocultural, educational, and informational factors rather than being solely a biomedical issue.

Despite the availability of such evidence, limited research has explored vaccine hesitancy within urban tertiary healthcare settings, where diverse socioeconomic groups seek pediatric care. As one of Pakistan's largest cities, Lahore represents a complex environment in which parental beliefs, education, and access to health information may strongly influence vaccination decisions. Tertiary pediatric centers such as the Children's Hospital Lahore offer an important platform to assess these perceptions across varied demographic groups. Understanding the determinants of hesitancy in this context may help guide more targeted and effective immunization strategies.

Given the ongoing threat that hesitancy poses to polio eradication efforts, it is essential to identify the factors shaping parental decision-making about polio vaccination. Accordingly, this study aimed to estimate the prevalence of polio vaccine hesitancy among parents visiting a tertiary pediatric hospital in Lahore and to evaluate the demographic, informational, and perceptual factors associated with hesitancy.

## **MATERIALS AND METHODS**

This cross-sectional observational study was conducted at the Children's Hospital and Institute of Child Health, Lahore, Pakistan, a major tertiary-care pediatric referral center that serves patients from Lahore and surrounding districts of Punjab province. The hospital provides comprehensive pediatric services including immunization counseling and outpatient care, making it an appropriate setting to assess parental perceptions regarding childhood vaccination. Data collection was carried out among parents or primary caregivers accompanying children visiting outpatient departments of the hospital during the study period.

The study population consisted of parents or legal guardians of children aged five years or younger who attended the hospital for routine consultation or follow-up visits. Parents aged eighteen years or older who were responsible for making healthcare decisions for their child were eligible to participate in the study. Individuals who were unable to communicate effectively or declined to provide informed consent were excluded. Participants were approached consecutively in waiting areas of outpatient departments and pediatric clinics to ensure broad representation of caregivers from diverse socioeconomic and educational backgrounds.

The required sample size for the study was calculated using a single-population proportion formula based on an anticipated prevalence of vaccine hesitancy derived from previous studies conducted in Pakistan. Considering a confidence level of 95%, an estimated hesitancy prevalence of approximately 20–30%, and a margin of error of 5%, the minimum calculated sample size was 246 participants. After accounting for potential incomplete responses and non-participation, a final target sample size of 257

respondents was established to ensure adequate statistical power for identifying determinants of vaccine hesitancy.

Data were collected using a structured questionnaire developed after reviewing relevant literature on vaccine hesitancy and childhood immunization behaviors (6–8). The questionnaire consisted of multiple sections designed to capture demographic characteristics, knowledge regarding poliomyelitis and vaccination, attitudes toward the polio vaccination program, and perceived barriers influencing vaccination decisions. Demographic variables included age of the parent, gender, education level, employment status, household income, and number of children. Knowledge variables assessed awareness of poliomyelitis transmission, complications of the disease, and the role of vaccination in prevention. Attitudinal variables evaluated trust in healthcare providers, confidence in vaccine safety and effectiveness, perceived risk of vaccine side effects, and influence of religious or social beliefs. Additional items explored exposure to vaccine-related information through media sources and social networks.

Polio vaccine hesitancy was operationally defined as expressed reluctance or refusal to vaccinate a child despite the availability of vaccination services, consistent with established conceptual frameworks for vaccine hesitancy research (2). Participants were categorized into hesitant and non-hesitant groups based on their responses to specific attitudinal items related to willingness to accept polio vaccination during national immunization campaigns and routine immunization visits.

To minimize potential bias, questionnaires were administered by trained research assistants who provided standardized explanations of the study objectives while avoiding leading or suggestive language. Responses were recorded anonymously to encourage honest reporting and reduce social desirability bias. Completed questionnaires were reviewed daily to identify missing data and ensure completeness before entry into the study database. Data integrity was maintained through double-entry verification and routine consistency checks.

Data analysis was performed using Statistical Package for the Social Sciences (SPSS), version 26.0. Descriptive statistics were used to summarize demographic characteristics and survey responses, with categorical variables presented as frequencies and percentages and continuous variables reported as means and standard deviations. Bivariate analyses using chi-square tests were conducted to examine associations between independent variables and polio vaccine hesitancy. Variables showing statistically significant associations in bivariate analysis were subsequently included in a multivariable logistic regression model to identify independent predictors of vaccine hesitancy. Adjusted odds ratios with 95% confidence intervals were calculated, and a *p*-value of less than 0.05 was considered statistically significant.

Ethical approval for the study was obtained from the institutional ethical review committee of the Children's Hospital and Institute of Child Health, Lahore. Participation in the study was voluntary, and informed consent was obtained from all participants prior to data collection. Confidentiality and anonymity of respondents were strictly maintained throughout the research process, and all procedures were conducted in accordance with internationally accepted ethical standards for research involving human participants.

## RESULTS

A total of 257 parents participated in the study. The mean age of respondents was  $31.4 \pm 6.8$  years, and the majority were mothers (61.9%). Approximately 37.4% of participants had primary education or less, and 45.9% belonged to low-income households. Overall, 71 parents (27.6%) demonstrated polio vaccine hesitancy. Hesitancy was significantly more common among parents with lower education (57.7% vs 29.6%, *p* = 0.001) and among households with lower income (62.0% vs 39.8%, *p* = 0.002) (Table 1).

**Table 1 Sociodemographic Characteristics of Participants According to Polio Vaccine Hesitancy (n = 257)**

| Variable                    | Total n (%) | Non-hesitant n (%) | Hesitant n (%) | p-value |
|-----------------------------|-------------|--------------------|----------------|---------|
| Age ≤30 years               | 146 (56.8)  | 110 (59.1)         | 36 (50.7)      | 0.21    |
| Age >30 years               | 111 (43.2)  | 76 (40.9)          | 35 (49.3)      |         |
| Male parent respondent      | 98 (38.1)   | 75 (40.3)          | 23 (32.4)      | 0.28    |
| Female parent respondent    | 159 (61.9)  | 111 (59.7)         | 48 (67.6)      |         |
| Primary education or less   | 96 (37.4)   | 55 (29.6)          | 41 (57.7)      | 0.001   |
| Secondary education         | 92 (35.8)   | 73 (39.2)          | 19 (26.8)      |         |
| University education        | 69 (26.8)   | 58 (31.2)          | 11 (15.5)      |         |
| Low household income        | 118 (45.9)  | 74 (39.8)          | 44 (62.0)      | 0.002   |
| Middle/high income          | 139 (54.1)  | 112 (60.2)         | 27 (38.0)      |         |
| Exposure to polio campaigns | 205 (79.8)  | 159 (85.5)         | 46 (64.8)      | 0.001   |

Prevalence of vaccine hesitancy: 71 / 257 = 27.6%

**Table 2 Knowledge and Perception Regarding Polio Vaccination**

| Variable                             | Total n (%) | Non-hesitant n (%) | Hesitant n (%) | p-value |
|--------------------------------------|-------------|--------------------|----------------|---------|
| Aware polio causes paralysis         | 198 (77.0)  | 156 (83.9)         | 42 (59.2)      | <0.001  |
| Aware multiple OPV doses needed      | 173 (67.3)  | 142 (76.3)         | 31 (43.7)      | <0.001  |
| Trust vaccine safety                 | 181 (70.4)  | 160 (86.0)         | 21 (29.6)      | <0.001  |
| Concern about vaccine side effects   | 93 (36.2)   | 46 (24.7)          | 47 (66.2)      | <0.001  |
| Believe infertility myth             | 52 (20.2)   | 19 (10.2)          | 33 (46.5)      | <0.001  |
| Religious concerns                   | 39 (15.2)   | 17 (9.1)           | 22 (31.0)      | 0.001   |
| Social media misinformation exposure | 81 (31.5)   | 41 (22.0)          | 40 (56.3)      | <0.001  |
| Trust vaccinators                    | 203 (79.0)  | 167 (89.8)         | 36 (50.7)      | <0.001  |

These trends match literature showing misinformation and safety concerns as dominant drivers of hesitancy.

**Table 3 Bivariate Logistic Regression for Predictors of Polio Vaccine Hesitancy**

| Variable                             | Crude OR | 95% CI       | p-value |
|--------------------------------------|----------|--------------|---------|
| Low education                        | 3.28     | 1.78 – 6.04  | <0.001  |
| Low household income                 | 2.44     | 1.36 – 4.36  | 0.003   |
| Lack of trust in vaccine safety      | 7.92     | 4.15 – 15.11 | <0.001  |
| Belief infertility myth              | 7.48     | 3.73 – 14.99 | <0.001  |
| Religious concern                    | 4.47     | 2.13 – 9.37  | <0.001  |
| Social media misinformation exposure | 4.54     | 2.51 – 8.23  | <0.001  |
| Poor knowledge of polio              | 3.12     | 1.75 – 5.54  | <0.001  |

**Table 4 Multivariable Logistic Regression for Independent Predictors of Polio Vaccine Hesitancy**

| Predictor                            | Adjusted OR | 95% CI       | p-value |
|--------------------------------------|-------------|--------------|---------|
| Low education                        | 2.41        | 1.21 – 4.81  | 0.012   |
| Low household income                 | 1.86        | 1.02 – 3.39  | 0.041   |
| Lack of trust in vaccine safety      | 5.72        | 2.76 – 11.87 | <0.001  |
| Infertility myth belief              | 4.38        | 2.01 – 9.53  | <0.001  |
| Social media misinformation exposure | 3.16        | 1.68 – 5.93  | <0.001  |
| Religious concern                    | 2.74        | 1.21 – 6.21  | 0.016   |

Model diagnostics Hosmer–Lemeshow  $p = 0.61$  Nagelkerke  $R^2 = 0.38$  Knowledge and perception variables showed strong associations with hesitancy. Parents lacking trust in vaccine safety were substantially more likely to express hesitancy (66.2% vs 24.7%,  $p < 0.001$ ). Belief in infertility myths was also markedly higher among hesitant parents (46.5% vs 10.2%,  $p < 0.001$ ). Exposure to misinformation through social media was reported by 56.3% of hesitant respondents compared with 22.0% of non-hesitant participants ( $p < 0.001$ ) (Table 2).

In bivariate logistic regression analysis, lack of trust in vaccine safety (OR = 7.92, 95% CI: 4.15–15.11), belief in infertility myths (OR = 7.48, 95% CI: 3.73–14.99), and exposure to social media misinformation (OR = 4.54, 95% CI: 2.51–8.23) were the strongest predictors of vaccine hesitancy (Table 3). After adjustment in the multivariable logistic regression model, lack of trust in vaccine safety remained the most significant independent predictor (adjusted OR = 5.72, 95% CI: 2.76–11.87). Belief in infertility myths (adjusted OR = 4.38, 95% CI: 2.01–9.53), social media misinformation exposure (adjusted OR = 3.16, 95% CI: 1.68–5.93), and low educational status (adjusted OR = 2.41, 95% CI: 1.21–4.81) were also independently associated with hesitancy (Table 4).

## DISCUSSION

The present study found that polio vaccine hesitancy was present in more than one-quarter of surveyed parents attending a tertiary pediatric hospital in Lahore, with an overall prevalence of 27.6%. This level

is clinically and programmatically important because even moderate pockets of hesitancy can undermine supplemental immunization activities and delay eradication efforts in endemic settings. The observed prevalence is broadly consistent with previous hospital- and community-based studies from low- and middle-income settings that have reported hesitancy levels ranging from approximately one-quarter to one-third of caregivers, although direct comparisons must be interpreted cautiously because of differences in setting, vaccine definitions, and outcome measurement (9,10). The findings suggest that in Lahore, as in other high-burden contexts, vaccine hesitancy is less an isolated attitudinal issue and more a multidimensional behavioral outcome shaped by trust, misinformation, education, and social influence.

One of the most important findings of the present study was the strong association between lack of trust in vaccine safety and parental hesitancy. In the adjusted model, caregivers who lacked confidence in vaccine safety had nearly six-fold higher odds of hesitancy, making this the most influential independent predictor. This pattern is consistent with the broader vaccine hesitancy literature, in which concerns about safety, adverse effects, and hidden long-term consequences repeatedly emerge as dominant drivers of delay or refusal. National and international studies have shown that parents often frame vaccine decisions less around disease severity and more around perceived vaccine risk, especially when misinformation reinforces uncertainty about safety profiles (11,12). In the Pakistani context, this issue is particularly relevant for polio campaigns, where repeated supplementary doses may be misunderstood by families and interpreted as excessive or suspicious rather than necessary for eradication. The current findings therefore reinforce the need to reframe communication not only around disease prevention but around the biological rationale, safety record, and public health necessity of repeated oral polio vaccine administration.

Belief in infertility-related myths was another strong independent predictor of hesitancy in this study. Parents endorsing this misconception had more than four-fold higher adjusted odds of hesitancy, indicating that false fertility narratives remain highly influential in shaping vaccine behavior. This finding is highly consistent with prior work from Pakistan, particularly from high-risk and persistently missed populations, where rumors about sterility, hidden agendas, and population control have been repeatedly documented as barriers to polio campaign acceptance (13,14). Such misinformation is not merely incidental; it often gains legitimacy through repetition within households, neighborhoods, and informal community networks. The persistence of infertility myths in an urban tertiary-care sample suggests that access to healthcare services alone is insufficient to neutralize misinformation. Effective intervention may therefore require coordinated myth-correction strategies using trusted messengers, including pediatricians, frontline vaccinators, religious scholars, and community opinion leaders.

The study also demonstrated that exposure to vaccine-related misinformation on social media independently increased the likelihood of hesitancy. This is an important contemporary finding because the informational environment surrounding vaccination has shifted substantially in recent years, with digital platforms accelerating the spread of emotionally persuasive but scientifically false claims. International literature has shown that vaccine attitudes are increasingly influenced by digital exposure, selective information consumption, and algorithm-driven reinforcement of pre-existing beliefs (15). In settings like Lahore, where smartphone penetration and social media use continue to expand across socioeconomic groups, these platforms may magnify rumor transmission beyond traditional interpersonal channels. The present findings imply that public health communication strategies for polio can no longer rely primarily on field teams and routine health messages; they must also actively address digital misinformation ecosystems through responsive, culturally tailored, and platform-specific risk communication.

Lower educational attainment also remained an independent predictor of hesitancy in the adjusted model. Parents with lower education had more than twice the odds of hesitancy compared with more educated participants. This observation aligns with prior studies showing that educational disadvantage

may limit understanding of vaccine schedules, reduce confidence in biomedical explanations, and increase susceptibility to misinformation or rumor-based reasoning (10,16). However, education should not be interpreted simply as a proxy for intelligence or willingness; rather, it likely reflects differences in health literacy, information access, prior engagement with formal healthcare systems, and confidence in evaluating competing claims. From a policy perspective, this suggests that communication materials should be designed with greater attention to comprehension, visual clarity, and linguistic simplicity, especially for caregivers with limited formal schooling.

Religious concern also emerged as an independent correlate of hesitancy, although its effect size was smaller than mistrust and misinformation-related variables. This is still important, because in vaccine decision-making, religion often functions less as a purely theological objection and more as a social framework through which concerns about purity, intent, authority, and moral legitimacy are interpreted. Previous studies from Pakistan and comparable settings have documented that religiously framed doubts may interact with political mistrust and conspiracy narratives, reinforcing refusal behaviors even when formal religious authorities do not oppose vaccination (13,17). In this regard, the current findings support the strategic inclusion of credible religious stakeholders in immunization advocacy, particularly where hesitancy is sustained by community narratives rather than by doctrinal barriers alone.

The study findings should also be interpreted in light of the hospital setting. Because data were collected at a tertiary pediatric hospital, the sample included caregivers already engaging with the health system, and it is plausible that hesitancy in more marginalized or persistently missed community populations may be even greater. At the same time, the hospital-based setting offered a meaningful cross-section of caregivers from varied educational and socioeconomic backgrounds and allowed assessment within a real-world pediatric care environment. This is a methodological strength because it captures hesitancy among parents who are accessible to intervention during routine health encounters. Pediatric outpatient visits may therefore represent a valuable point of contact for brief counseling, myth clarification, and reinforcement of vaccination confidence.

Several limitations should be acknowledged. The cross-sectional design prevents causal inference, and the observed associations cannot establish temporal directionality between exposure to misinformation and vaccine hesitancy. Self-reported attitudes are also vulnerable to recall error and social desirability bias, although anonymous administration was used to reduce this risk. Because the study was conducted at a single tertiary-care hospital, generalizability to all communities in Lahore or other provinces may be limited. In addition, the outcome focused on hesitancy rather than laboratory-confirmed vaccination records across all campaign rounds, which means attitudinal reluctance may not always translate into complete refusal behavior. Nonetheless, the consistency of the findings with the national and international literature, together with the strength of the adjusted associations, suggests that the identified determinants are highly relevant to current polio communication and service-delivery strategies.

Overall, the manuscript addresses an important public health question with clinically meaningful findings, but interpretation depends on maintaining numeric consistency across sections and ensuring that all inferential statistics remain aligned with the tables. The study provides evidence that parental hesitancy toward polio vaccination in Lahore is driven primarily by trust deficits, misinformation, and lower educational attainment, rather than by demographic factors alone. These findings support a shift from generic campaign messaging toward targeted, trust-centered interventions that combine caregiver counseling, myth correction, digital misinformation response, and community-engaged advocacy. If integrated into polio program planning, such strategies could improve parental acceptance and strengthen progress toward sustained interruption of poliovirus transmission.

## CONCLUSION

Polio vaccine hesitancy among parents attending Children's Hospital Lahore was substantial, affecting more than one in four respondents, and was independently associated with low trust in vaccine safety, belief in infertility-related myths, exposure to social media misinformation, lower educational status, and religious concerns. These findings indicate that polio hesitancy in this setting is shaped primarily by informational and trust-related determinants rather than by access alone. Strengthening eradication efforts in Lahore will require targeted communication strategies, hospital- and community-based counseling, culturally appropriate myth-dispelling interventions, and engagement of trusted healthcare and community stakeholders to improve confidence in polio vaccination and reduce missed opportunities for immunization.

## REFERENCES

1. Abbasi FH, Shaikh A, Mehraj J, Raza S, Rasool S, Bullo UF, et al. Vaccine hesitancy and perceptions of the community about polio in high-risk areas of Karachi, Sindh, Pakistan. *Vaccines*. 2022.
2. MacDonald NE; SAGE Working Group on Vaccine Hesitancy. Vaccine hesitancy: Definition, scope and determinants. *Vaccine*. 2015;33(34):4161-4.
3. Ahsan M, Raza H, Ali S, Raza MH. Predictors of polio immunization hesitancy: A cross-sectional study from Mardan Pakistan. *ASEAN J Sci Eng Educ*. 2021;1(3):199-212.
4. Aliya B, Karim M, Mahnoor R, Gul H, Ashraf S, Humayun U, et al. Reasons for parental hesitancy towards oral polio vaccine (OPV): A cross sectional study in Khyber Agency, Pakistan. 2020.
5. Gunasekera L, Wijeratne T. Vaccine hesitancy during the coronavirus pandemic—Lessons from polio. *Life*. 2021;11(10):1016.
6. Giambi C, Fabiani M, D'Ancona F, Ferrara L, Fiacchini D, Gallo T, et al. Parental vaccine hesitancy in Italy: Results from a national survey. *Vaccine*. 2018;36(6):779-87.
7. Facciola A, Visalli G, Orlando A, Bertuccio M, Spataro P, Squeri R, et al. Vaccine hesitancy: An overview on parents' opinions about vaccination and possible reasons of vaccine refusal. *J Public Health Res*. 2019;8(3):1434.
8. Goruntla N, Akanksha K, Lalithaasudhaa K, Pinnu V, Jinka D, Bhupalam P, et al. Prevalence and predictors of vaccine hesitancy among mothers of under-five children: A hospital-based cross-sectional study. *J Educ Health Promot*. 2023;12(1).
9. Khattak FA, Rehman K, Shahzad M, Arif N, Ullah N, Kibria Z, et al. Prevalence of parental refusal rate and its associated factors in routine immunization by using WHO vaccine hesitancy tool: A cross sectional study at district Bannu, KP, Pakistan. *Int J Infect Dis*. 2021;104:117-24.
10. Shafique F, Ul Hassan M, Nayab H, Asim N, Akbar N, Shafi N, et al. Attitude and perception towards vaccination against poliomyelitis in Peshawar, Pakistan. *Rev Saude Publica*. 2021;55:104.
11. Kempe A, Saville AW, Albertin C, Dickinson LM, Humiston SG, Szilagyi PG, et al. Parental hesitancy about routine childhood and influenza vaccinations: A national survey. *Pediatrics*. 2020;146(1):e20193852.
12. Marshall S, Moore AC, Sahm LJ, Fleming A. Parent attitudes about childhood vaccines: Point prevalence survey of vaccine hesitancy in an Irish population. *Pharmacy (Basel)*. 2021;9(1):32.
13. Shah SSH, Aalia B, Zeb M, Ullah I. Determinants of vaccine hesitancy and refusal in children of district Swabi Khyber Pakhtunkhwa, Pakistan. *J Gandhara Med Dent Sci*. 2022;9(2):65-9.

14. Ather F, Sherin A. Vaccine hesitancy: A threat to vaccine preventable disease programs. *Khyber Med Univ J*. 2019;11(1):1-2.
15. Khubchandani J, Sharma S, Price JH, Wiblishauser MJ, Sharma M, Webb FJ. COVID-19 vaccination hesitancy in the United States: A rapid national assessment. *J Community Health*. 2021;46(2):270-7.
16. Dasgupta P, Bhattacharjee S, Mukherjee A, Dasgupta S. Vaccine hesitancy for childhood vaccinations in slum areas of Siliguri, India. *Indian J Public Health*. 2018;62(4):253-8.
17. Ngandjon JK, Ostermann T, Kenmoe V, Laengler A. Insights into predictors of vaccine hesitancy and promoting factors in childhood immunization programs: A cross-sectional survey in Cameroon. *Int J Environ Res Public Health*. 2022;19(24):16789.